arXiv:1001.1646v2 [math.NT] 14 Jan 2010

THE CATENARY AND TAME DEGREE OF NUMERICAL MONOIDS GENERATED BY GENERALIZED ARITHMETIC SEQUENCES

M. OMIDALI

Abstract. Studying ceratin combinatorial properties of non-unique factorizations have been a subject of recent literatures. Little is known about two combinatorial invariants, namely the catenary degree and the tame degree, even in the case of numerical monoids. In this paper we compute these invariants for a certain class of numerical monoids generated by generalized arithmetic sequences. We also show that the difference between the tame degree and the catenary degree can be arbitrary large even if the number of minimal generators is fixed.

Keywords. Numerical monoids, Tame degree, Catenary degree
2010 Mathematics Subject Classification. 20M13

1. Introduction

Various aspects of non-unique factorizations in integral domains have been a subject of researchers in recent years. There are several arithmetic invariants that measure the behavior of non-unique factorizations in integral domains. Early works on the behavior of non-unique factorizations in integral domains were focused on the length of irreducible factorizations of an element. Delta sets and the elasticity are some of these invariants that measure how far an integral domain or monoid is from being factorial or half-factorial (i.e., all the irreducible factorizations of an element have the same length). In recent years other invariants which have close relations with the distance between irreducible factorizations have been appearing in literatures. The *catenary degree* and the *tame degree* are such invariants. The monograph of Geroldinger and Halter-Koch [10], the recent survey [11] and [9] are good references for studying the catenary degree and the tame degree in commutative cancellative monoids.

Throughout $\mathbb{N}$ will be the set of non-negative integers. A numerical monoid is a submonoid of $\mathbb{N}$ (that is closed under addition) which contains 0 and its complement in $\mathbb{N}$ is finite. Every numerical monoid is necessarily finitely generated and has a minimal set of generators. For a set $\{n_1, \ldots, n_p\}$ of increasing positive integers, the numerical monoid generated by n_i's is $\langle n_1, \ldots, n_p \rangle := \{\sum_{i=1}^{p} z_i n_i | z_i \in \mathbb{N},\ i = 1, \ldots, p\}$. Delta sets of numerical monoids were studied in [1, 3], and it is shown that in numerical monoids generated by arithmetic sequences, Delta sets reduce to a single element. This element is the difference between two consecutive elements of the arithmetic sequence. Also in [7] it is shown the elasticity of a numerical monoid equals the quotient of the largest by the smallest minimal generator. For numerical monoids generated by an arithmetic sequence the catenary degree and the tame degree were determined in [4].

In this paper we consider numerical monoids generated by generalized arithmetic sequences, that is numerical monoids of the form $S = \langle a, ha + d, \ldots, ha + xd \rangle$, where a, h, x, and d are positive integers and $\gcd(a, d) = 1$. This class of numerical monoids contains all numerical monoids generated by arithmetic sequences. We explicitly compute their catenary and tame degrees. As a result we show that the difference between the tame degree and the catenary degree can be arbitrary large even if the number of minimal generators is fixed. From [[10], Example 3.1.6] it is known that

$$\mathrm{c}(S) \leq \mathrm{t}(S) \leq \frac{g(S) + n_p}{n_1} + 1,$$

where $\mathrm{c}(S)$ and $\mathrm{t}(S)$ are the catenary degree and the tame degree of S and $g(S)$ is the Frobenius number of S (i.e. the largest positive integer not belonging to S). For numerical monoids generated by generalized arithmetic sequences we show that the difference between $\mathrm{c}(S)$ and $(g(S) + n_p)/n_1 + 1$ can be arbitrarily large. Moreover, $\mathrm{t}(S)$ can be $\mathrm{c}(S)$ or $\lfloor (g(S) + n_p)/n_1 + 1 \rfloor$ and all of these things can happen even if the number of minimal generators is fixed.

For general theory of numerical monoids we refer the reader to [15]. Other aspects of numerical monoids generated by generalized arithmetic sequences are studied in [2], [13], [14], [16], [17]. We mainly use a membership criterion presented in [14]. Computing some combinatorial invariants of numerical monoids, or more generally commutative cancellative monoids such as the elesticity and the tame degree requires us to determine the set of irreducible elements of a congruence related to the monoid [6]. All the results in this paper have been tested with the Numericalsgps package of GAP [8].

Notation 1.1. For a rational number r by $\lceil r \rceil$ we mean the least integer bigger than or equal to r and by $\lfloor r \rfloor$ we mean the greatest integer less than or equal to r. Also $(m \mod n)$ means the reminder of quotient of the integer division of m by n where $m, n \in \mathbb{Z}$. For integers a, b, and c by $a \equiv b (\mathrm{mod} c)$ we mean that $a - b$ is divisible by c.

2. Preliminaries

Suppose that S is a numerical monoid which is minimally generated by $\{n_1, \ldots, n_p\}$. Each n_i is called an atom and $\{n_1, \ldots, n_p\}$ is called the set of atoms of S and is denoted by $\mathcal{A}(S)$. Define a partial order $\leq_S$ on S by declaring $a \leq_S b$ if $b - a \in S$. The factorization morphism of S is

$$\varphi : \mathbb{N}^p \longrightarrow S, \ \ \varphi(z_1, \ldots, z_p) = z_1 n_1 + \ldots + z_p n_p.$$

Then S is isomorphic to $\mathbb{N}^p/\sigma$, where $a\sigma b$ if $\varphi(a) = \varphi(b)$. The set of factorizations of $n \in S$ is defined by

$$\mathsf{Z}(n) = \varphi^{-1}(n) = \{(z_1, \ldots, z_p) \in \mathbb{N}^p | z_1 n_1 + \ldots + z_p n_p = n\}.$$

If $z = (z_1, \ldots, z_p)$ is a factorization of n, then the length of z is $|z| = z_1 + \ldots + z_p$. The set of lengths of n is defined by

$$\mathsf{L}(n) = \{|z| \bigm| z \in \mathsf{Z}(n)\} \subset \mathbb{N}_0.$$

Also the support of z is defined by

$$\mathrm{supp}(z) = \{i \in \{1, \ldots, p\} | z_i \neq 0\}.$$

For $z = (z_1, \ldots, z_p), z' = (z'_1, \ldots, z'_p) \in \mathbb{N}^p$ we set

$$\gcd(z, z') = (\min\{z_1, z'_1\}, \ldots, \min\{z_p, z'_p\}), \frac{z}{z'} = z - z'.$$

The distance between z and z' is defined by

$$\mathrm{d}(z, z') = \max\Big\{\Big|\frac{z}{\gcd(z, z')}\Big|, \Big|\frac{z'}{\gcd(z, z')}\Big|\Big\}.$$

For two nonempty subsets X and Y of $\mathbb{N}^p$ we define the distance between X and Y as follows

$$\mathrm{d}(X, Y) = \min\{\mathrm{d}(x, y) | x \in X, y \in Y\}.$$

If $X = \{x\}$ then we use $\mathrm{d}(x, Y)$ instead of $\mathrm{d}(X, Y)$.

Let $n \in S$ and $z, z' \in \mathsf{Z}(n)$. Then an $N-$*chain of factorizations* from z to z' is a sequence $z_0, \ldots, z_k \in \mathsf{Z}(n)$ such that $z = z_0$, $z' = z_k$ and $\mathrm{d}(z_i, z_{i+1}) \leq N$ for all i. The catenary degree of n, $\mathrm{c}(n)$, is defined as the least non-negative integer N such that for any two factorizations $z, z' \in \mathsf{Z}(n)$, there is an N-chain from z to z'. The catenary degree of S is defined by

$$\mathrm{c}(S) = \sup\{\mathrm{c}(n) | n \in S\}.$$

Two elements z and z' of $\mathbb{N}^p$ are $\mathscr{R}$-related if there exists a sequence $z = z_0, \ldots, z_k = z'$ in $\mathbb{N}^p$ such that $\mathrm{supp}(z_i) \cap \mathrm{supp}(z_{i+1}) \neq \varnothing$, for every $i \in \{1, \ldots, k-1\}$. As the number of different factorizations of an element $n \in S$ is finite, so is the number of different $\mathscr{R}$-classes of $\mathsf{Z}(n)$. Let $n \in S$ and let $\mathscr{R}_1^n, \ldots, \mathscr{R}_{k_n}^n$ be the different $\mathscr{R}$-classes of elements in $\mathsf{Z}(n)$. Set $\mu(n) = \max\{r_1^n, \ldots, r_{k_n}^n\}$, where $r_i^n = \min\{|z| \big| z \in \mathscr{R}_i^n\}$. Define

$$\mu(S) = \max\{\mu(n) \big| n \in S \text{ and } k_n \geq 2\}.$$

Theorem 2.1 ([5], Theorem 3.1)**.** *Let S be a numerical monoid. Then*

$$\mathrm{c}(S) = \mu(S).$$

Let $n \in S$. For $i \in \{1, \ldots, p\}$ we set $\mathsf{Z}^i(n) = \{(z_1, \ldots, z_p) \in \mathsf{Z}(n) | z_i \neq 0\}$. Suppose that $n - n_i \in S$ for some $i \in \{1, \ldots, p\}$. We define

$$\mathrm{t}_i(n) = \max\{\mathrm{d}(z, \mathsf{Z}^i(n)) | z \in \mathsf{Z}(n)\}.$$

Setting

$$\mathrm{t}(n) = \max\{\mathrm{t}_i(n) | n - n_i \in S, 1 \leq i \leq p\},$$

and

$$\mathrm{t}_i(S) = \max\{\mathrm{t}_i(n) | n - n_i \in S\}, \quad i = 1, 2, \ldots, p,$$

we define the tame degree of S by

$$\mathrm{t}(S) = \max\{\mathrm{t}(n) | n \in S\} = \max\{\mathrm{t}_i(S) | i = 1, 2, \ldots, p\}.$$

It is clear from definitions that $\mathrm{c}(n) \leq \mathrm{t}(n) \leq \max \mathsf{L}(n)$.

For $a, b \in S$ let $\omega(a, b)$ be the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ with the following property:

- For all $n \in \mathbb{N}$ and $a_1, \ldots, a_n \in S$, if $a = a_1 + \ldots + a_n$ and $b \leq_S a$ then there exists a subset $\Omega \subset [1, n]$ such that $|\Omega| \leq N$ and
$$b \leq_S \sum_{\nu \in \Omega} a_\nu.$$

Note that if $b \nleq_S a$, then set $\omega(a, b) = 0$. For $b \in S$ we define

$$\omega(S, b) = \sup\{\omega(a, b) \mid a \in S\} \in \mathbb{N}_0 \cup \{\infty\}.$$

For $k \in \mathbb{N}$ and $b \in S$ we set

$$\begin{aligned}\tau_k(S, b) = &\sup\{\min \mathsf{L}(a - b) \big| a = u_1 + \ldots + u_j \in b + S \text{ with } j \in [0, k],\\ &u_1, \ldots, u_j \in \mathcal{A}(S), \text{and } b \nleq_S a - u_i \text{ for all } i \in [1, j]\} \in \mathbb{N}_0 \cup \{\infty\}\end{aligned}$$

and

$$\tau(S, b) = \sup\{\tau_k(S, b) \big| k \in \mathbb{N}\} \in \mathbb{N}_0 \cup \{\infty\}.$$

We define the ω invariant and the τ invariant of S as

$$\omega(S) = \sup\{\omega(S, n_i) \mid 1 \leq i \leq p\} \in \mathbb{N}_0 \cup \{\infty\},$$

$$\tau(S) = \sup\{\tau(S, n_i) \mid 1 \leq n_i \leq p\} \in \mathbb{N}_0 \cup \{\infty\}.$$

Theorem 2.2. *If $S \neq \mathbb{N}$, then for every $1 \leq i \leq p$ we have*

$$\mathrm{t}_i(S) = \max\{\omega(S, n_i), 1 + \tau(S, n_i)\} \in \mathbb{N}_{\geq 2} \cup \{\infty\}.$$

Proof. Since $S \neq \mathbb{N}$, all n_i are atoms but not primes. Thus the assertion follows from [[12], Theorem 3.6]. $\square$

3. Numerical monoid generated by generalized arithmetic sequences

Let $S = \langle a, ha+d, \dots, ha+xd\rangle$ be a numerical monoid where a, d, h, and x are positive integers and $\gcd(a, d) = 1$. If $i \geq a$ then $ha + id = \lfloor \frac{i}{a} \rfloor da + (ha + (i \mod a)d) \in S$. Therefore we may assume that $x < a$. In this case $\{a, ha + d, \dots, ha + xd\}$ is the minimal set of generators of S ([14], Corollary 2.5). We sometimes use $n_0, n_1, \dots, n_x$ instead of $a, ha + d, \dots, ha + xd$, respectively.

Theorem 3.1 ([14], Proposition 2.1)**.** *Suppose that $S = \langle a, ha + d, \dots, ha+xd\rangle$ where a, d, h, and x are positive integers,* $\gcd(a, d) = 1$*, and $1 \leq x \leq a - 1$. Let $n = qa + id$ where $q, i \in \mathbb{N}$ and $0 \leq i \leq a - 1$. Then $n \in S$ if and only if $\lceil \frac{i}{x} \rceil h \leq q$.*

Note that by a generalization of the euclidean algorithm every integer n has a unique representation of the form $n = qa + id$ with $0 \leq i \leq a - 1$. So the above condition gives a simple membership criterion for numerical monoids generated by generalized arithmetic sequences.

Let S be a numerical monoid and $n \in S$. The Apéry set associated to n is defined as

$$\mathrm{Ap}(S, n) = \{s \in S | s - n \notin S\}.$$

It is easy to see that $\mathrm{Ap}(S, n)$ has n elements of different congruence classes modulo n. Therefore each element of S can be uniquely expressed as $kn + w$, where $k \in \mathbb{N}$ and $w \in \mathrm{Ap}(S, n)$.

Theorem 3.2 ([14], Proposition 2.6)**.** *Let $S = \langle a, ha+d, \dots, ha+xd\rangle$ where a, d, h, and x are positive integers,* $\gcd(a, d) = 1$*, and $1 \leq x \leq a - 1$. Then*

$$\mathrm{Ap}(S, a) = \{\lceil \frac{i}{x} \rceil ha + id | 0 \leq i \leq a - 1\}.$$

Let S be a numerical monoid generated minimally by $n_1, n_2, \dots, n_p$. For every $n \in S$ we define

$$C_n = \{V \subseteq \{1, 2, \dots, p\} | \exists z \in \mathsf{Z}(n), V \subseteq \mathrm{supp}(z)\}.$$

It is easy to see that C_n is a simplicial complex and the number of $\mathscr{R}$-classes of n is equal to the number of path components of C_n. We define a graph $G_n = (V_n, E_n)$ as

$$V_n = \{i \in \{1, 2, \ldots, p\} | n - n_i \in S\},$$
$$E_n = \{\{i, j\} | n - (n_i + n + j) \in S, i \neq j \in \{1, 2, \ldots, p\}\},$$

with V_n and E_n are the sets of vertices and edges of G_n, respectively. Then the number of path components of C_n and G_n are equal and thus the number of $\mathscr{R}$-classes of n is equal to the number of path components of G_n.

Example. Let $S = \langle 5, 6, 7, 9 \rangle$ and $n = 18$. Then 18 can be written in terms of minimal generators as follows:

$$18 = 2 \times 9 = 5 + 6 + 7.$$

C_{18} and G_{18} are depicted in Figure 1.



C_{18} G_{18}

FIGURE 1. C_n and G_n for $S = \langle 5, 6, 7, 9 \rangle$ and $n = 18$

In the proof of [[14], Theorem 2.16] we found all elements n of a numerical monoid generated by a generalized arithmetic sequence for which G_n is disconnected, and we summarize it in the following Theorem.

Theorem 3.3. *Suppose that* $S = \langle a, ha + d, \ldots, ha + xd \rangle$ *where* $a, d, h,$ *and* x *are positive integers,* $\gcd(a, d) = 1$*, and* $1 \leq x \leq a - 1$*. Let* $n \in S$ *be such that* C_n *is disconnected. Then either* $n = n_i + n_j$*, with* $1 \leq i, j \leq x$*, or* $n \in \{pn_x + n_{r+1}, \ldots, pn_x + n_x = (p + 1)n_x\}$*, where* $p = \lfloor \frac{a-1}{x} \rfloor$ *and* $r = (a - 1) \mod x$*.*

Proposition 3.4. *Suppose that* $S = \langle a, ha + d, \ldots, ha + xd \rangle$ *where* $a, d, h,$ *and* x *are positive integers,* $\gcd(a, d) = 1$*, and* $1 \leq x \leq a - 1$*. Let* $n = qa + id \in S$ *be such that* $0 \leq i < a$*. Then*

$$\max \mathsf{L}(n) = q - \lceil \frac{i}{x} \rceil (h - 1).$$

Proof. First we show that $\max \mathsf{L}(n) \geq q - \lceil \frac{i}{x} \rceil (h - 1)$. Let $i = \lceil \frac{i}{x} \rceil x - s$ with $0 \leq s < x$.

- If $s = 0$ then

$$n = qa + id = qa + \lceil \frac{i}{x} \rceil xd = (q - \lceil \frac{i}{x} \rceil h)a + \lceil \frac{i}{x} \rceil (ha + xd).$$

Therefore

$$\max \mathsf{L}(n) \geq (q - \lceil \frac{i}{x} \rceil h) + \lceil \frac{i}{x} \rceil = q - \lceil \frac{i}{x} \rceil (h-1).$$

- If $s > 0$ then $1 \leq \lceil \frac{i}{x} \rceil$ and $\lceil \frac{i}{x} \rceil h \leq q$. Therefore

$$\begin{aligned} n &= qa + id \\ &= qa + (\lceil \frac{i}{x} \rceil x - s)d \\ &= (q - \lceil \frac{i}{x} \rceil h)a + (\lceil \frac{i}{x} \rceil - 1)(ha + xd) + (ha + (x-s)d). \end{aligned}$$

Thus

$$\max \mathsf{L}(n) \geq (q - \lceil \frac{i}{x} \rceil h) + (\lceil \frac{i}{x} \rceil - 1) + 1 = q - \lceil \frac{i}{x} \rceil (h-1).$$

Now we show that $\max \mathsf{L}(n) \leq q - \lceil \frac{i}{x} \rceil (h-1)$. We use induction on q to show this. If $q = 0$ then $\lceil \frac{i}{x} \rceil h \leq 0$. Therefore $i = 0$, $n = 0$, and $0 = \max \mathsf{L}(n) \leq q - \lceil \frac{i}{x} \rceil (h-1)$. Now let $q > 0$. We have

$$\max \mathsf{L}(n) = 1 + \max\{\max \mathsf{L}(n') | n' \in \{n - n_0, n - n_1, \ldots, n - n_x\} \cap S\}.$$

So it is enough to show that for any $0 \leq j \leq x$, if $n' = n - n_j \in S$, then $\max \mathsf{L}(n') + 1 \leq q - \lceil \frac{i}{x} \rceil (h-1)$. There are three cases.

- First let $j = 0$ and $n' = n - n_0 = (q-1)a + id \in S$. Then by induction

$$\max \mathsf{L}(n') + 1 \leq (q-1) - \lceil \frac{i}{x} \rceil (h-1) + 1 = q - \lceil \frac{i}{x} \rceil (h-1).$$

- Now let $1 \leq j \leq i$ and $n' = n - n_j = (q-h)a + (i-j)d \in S$. Then $0 \leq i - j < a$ and $\lceil \frac{i-j}{x} \rceil \geq \lceil \frac{i-x}{x} \rceil = \lceil \frac{i}{x} \rceil - 1$. By induction we have

$$\max \mathsf{L}(n') + 1 \leq (q-h) - \lceil \frac{i-j}{x} \rceil (h-1) + 1 \leq q - \lceil \frac{i}{x} \rceil (h-1).$$

- Finally suppose that $i < j$ and $n' = n - n_j = (q-h)a + (i-j)d \in S$. Then $n' = (q-h-d)a + (i-j+a)d$ with $0 \leq i-j+a < a$. By induction and the fact that $i - j + a \geq i$ we have

$$\max \mathsf{L}(n') + 1 \leq (q-h-d) - \lceil \frac{i-j+a}{x} \rceil (h-1) + 1 \leq q - \lceil \frac{i}{x} \rceil (h-1).$$

□

Corollary 3.5. *Suppose that* $S = \langle a, ha+d, \ldots, ha+xd\rangle$ *where* $a, d, h,$ *and* x *are positive integers,* $\gcd(a, d) = 1$*, and* $1 \leq x \leq a-1$*. Let* $n = qa + id$ *then* $n \in S$ *if and only if*

$$\lceil \frac{i \mod a}{x} \rceil h \leq q + \lfloor \frac{i}{a} \rfloor d,$$

and in this case

$$\max \mathsf{L}(n) = (q + \lfloor \frac{i}{a} \rfloor d) - \lceil \frac{i \mod a}{x} \rceil (h-1).$$

Proof. Just note that $n = (q + \lfloor \frac{i}{a} \rfloor d)a + (i \mod a)d$. □

3.1. **The catenary degree.**

We are ready to find the catenary degree of numerical monoids generated by generalized arithmetic sequences.

Theorem 3.6 ([4], Proposition 5)**.** *Let* S *be a numerical monoid minimally generated by* $\{n_1, \ldots, n_p\}$ *with* $0 < n_1 < \ldots < n_p$*. Then*

$$\min\{k \in \mathbb{N} \setminus \{0\} \mid kn_1 \in \langle n_2, \ldots, n_p\rangle\} \leq \mathrm{c}(S). \tag{1}$$

Even though there are numerical monoids for which the inequality (1) is strict, we will show that it is an equality for numerical monoids generated by generalized arithmetic sequences. Not all numerical monoids benefit from this nice property. It even fails for numerical monoids generated by almost arithmetic sequences (i.e., numerical monoids generated by a set for which all but one element form some consecutive elements of an arithmetic sequence). For example any three relatively prime integers form an almost arithmetic sequence but $S = \langle 6, 9, 11\rangle$ for which we have $\mathrm{c}(S) = 4$ but $\min\{k \in \mathbb{N} \setminus \{0\} | 6k \in \langle 9, 11\rangle\} = 3$.

Lemma 3.7. *Let* $a, h, x,$ *and* d *be positive integers with* $x < a$ *and* $\gcd(a, d) = 1$*. Then*

$$\min\{k \in \mathbb{N} \setminus \{0\} | \, ka \in \langle ha+d, \ldots, ha+xd\rangle\} = \lceil \frac{a}{x} \rceil h + d.$$

Proof. Let $c = \gcd(ha+d, \ldots, ha+xd) = \gcd(h, d)$. Then

$$\min\{k \in \mathbb{N} \setminus \{0\} \mid ka \in \langle ha+d, \ldots, ha+xd\rangle\}$$
$$= c \min\{k \in \mathbb{N} \setminus \{0\} \mid ka \in \langle \frac{h}{c}a + \frac{d}{c}, \ldots, \frac{h}{c}a + x\frac{d}{c}\rangle\}.$$

Therefore we must show that

$$\min\{k \in \mathbb{N} \setminus \{0\} \mid ka \in \langle \frac{h}{c}a + \frac{d}{c}, \ldots, \frac{h}{c}a + x\frac{d}{c}\rangle\} = \lceil \frac{a}{x} \rceil \frac{h}{c} + \frac{d}{c}.$$

So we may assume that $\gcd(h, d) = 1$. Let $a' = ha+d, h' = 1, x' = x-1$ and $d' = d$. Then $S' = \langle a', h'a'+d', \ldots, h'a'+x'd'\rangle = \langle ha+d, \ldots, ha+$

$xd\rangle$ is a numerical monoid generated by a generalized arithmetic sequence. Let $q'a' + i'd = ka$ with $0 \le i' < a'$ and $\lceil \frac{i'}{x'} \rceil h' \le q'$ and $k \ne 0$. $\lceil \frac{i'}{x'} \rceil h' \le q'$ is equivalent to $i' \le q'x'$. Since $k \ne 0$ we have $q' > 0$. From $q'a' + i'd = ka$ we have $(q' + i')d = (k - q'h)a$ and from $\gcd(a, d) = 1$ we deduce that there is $j > 0$ such that $q' + i' = ja$ and $k - q'h = jd$. From $q' + i' = ja$ and $i' \le q'x'$ we deduce that $ja - q' \le q'x'$ or equivalently $ja \le q'(x' + 1) = q'x$. Thus, if $k > 0$, then $ka \in S'$ if and only if there are $q' > 0$ and $j > 0$ such that

$$ja \le q'x, \text{ and } k = q'h + jd.$$

The minimum of k with the stated property is attained if we set $j = 1$. In this case $a \le q'x$ and thus $\lceil \frac{a}{x} \rceil \le q'$. So the minimum of $k > 0$ with the property $ka \in S'$ is when $q' = \lceil \frac{a}{x} \rceil$ and $j = 1$ which is $k = q'h + d = \lceil \frac{a}{x} \rceil h + d$. □

Theorem 3.8. *Let* $S = \langle a, ha + d, \ldots, ha + xd\rangle$ *where* $a, d, h,$ *and* x *are positive integers,* $\gcd(a, d) = 1$, *and* $1 \le x \le a - 1$. *Then*

$$\mathrm{c}(S) = \lceil \frac{a}{x} \rceil h + d.$$

Proof. We know that $\mathrm{c}(S) \ge \lceil \frac{a}{x} \rceil h + d$ by Theorem 3.6 and Lemma 3.7. We complete the proof by showing that if $n \in S$ is such that the number of $\mathscr{R}$-classes of n is bigger than one then $\max \mathsf{L}(n) \le \lceil \frac{a}{x} \rceil h + d$. By Theorem 3.3 we have two cases.

- First let $n = n_i + n_j = 2ha + (i + j)d$ with $1 \le i, j \le x$. Then

$$\begin{aligned} \mathrm{c}(n) \le \max \mathsf{L}(n) &= 2h + \lfloor \frac{i+j}{a} \rfloor d - \lceil \frac{(i+j) \bmod a}{x} \rceil (h-1) \\ &\le 2h + \lfloor \frac{i+j}{a} \rfloor d \\ &\le 2h + d \\ &\le \lceil \frac{a}{x} \rceil h + d. \end{aligned}$$

- Now let $n \in \{pn_x + n_{r+1}, \ldots, pn_x + n_x = (p + 1)n_x\}$, with $p = \lfloor \frac{a-1}{x} \rfloor$ and $r = (a - 1) \bmod x$. If $r + 1 \le j < x$ and $n = pn_x + n_j$ then $n = (\lfloor \frac{a-1}{x} \rfloor + 1)ha + jd = \lceil \frac{a}{x} \rceil ha + jd$ and

$$\max \mathsf{L}(n) = \lceil \frac{a}{x} \rceil h - \lceil \frac{j}{x} \rceil (h-1) \le \lceil \frac{a}{x} \rceil h + d.$$

  If $n = (p + 1)n_x$ then $n = \lceil \frac{a}{x} \rceil ha + (px + x)d = (\lceil \frac{a}{x} \rceil h + d)a + (px + x - a)d = (\lceil \frac{a}{x} \rceil h + d)a + (x - 1 - r)d$. Since $0 \le x - 1 - r < x$ we have

$$\max \mathsf{L}(n) = \lceil \frac{a}{x} \rceil h + d - \lceil \frac{x-1-r}{x} \rceil (h-1) \le \lceil \frac{a}{x} \rceil h + d.$$

□

3.2. **The tame degree.** Now we find the tame degree of numerical monoids generated by generalized arithmetic sequences.

Theorem 3.9 ([4], Theorem 16). *Let S be a numerical monoid minimally generated by $\{n_1 < \ldots, n_p\}$. Let $n \in S$ be minimal satisfying $\mathrm{t}(n) = \mathrm{t}(S)$. Then $n = w + n_i$ for some $i \in \{1, \ldots, p\}$ and $w \in \mathrm{Ap}(S, n_j)$ with $j \in \{1, \ldots, p\} \setminus \{i\}$.*

Theorem 3.10. *Let $S = \langle a, ha + d, \ldots, ha + xd\rangle$ where $a, d, h,$ and x are positive integers, $\gcd(a, d) = 1$, and $1 \le x \le a - 1$. Then*

$$\mathrm{t}(S) = \omega(S) = \big(\lceil\frac{a-1}{x}\rceil + 1\big)h + d.$$

Proof. We set $t = \lceil\frac{a-1}{x}\rceil h + h + d$ and $n = ta$. Then

$$w = n - n_1 = (\lceil\frac{a-1}{x}\rceil h + d)a - d = \lceil\frac{a-1}{x}\rceil ha + (a-1)d \in \mathrm{Ap}(S, a)$$

Therefore there exists a factorization $z \in \mathsf{Z}^1(n)$. Also $z' = (t, 0, \ldots, 0) \in \mathsf{Z}(n) \setminus \mathsf{Z}^1(n)$ and therefore $\mathrm{t}(S) \ge \mathrm{t}_1(n) \ge \mathrm{d}(z', \mathsf{Z}^1(n)) = |z'| = t$. We complete the proof by concluding that $\mathrm{t}(S) \le t$. Let $n \in S$ be minimal such that $\mathrm{t}(n) = \mathrm{t}(S)$. Assume that z is a factorization of n such that $\mathrm{d}(z, z') = \mathrm{t}(n)$ with $z' \in \mathsf{Z}^i(n)$. From the minimality of n, we can derive that $\mathrm{supp}(z) \cap \mathrm{supp}(z') = \varnothing$ and thus $\mathrm{d}(z, z') = \max\{|z|, |z'|\}$. Let j be in the support of z. Theorem 3.9 ensures that $n - (n_i + n_j) \notin S$. Thus $n = (n - n_i) + n_i = (n - n_j) + n_j$ with $n - n_j \in \mathrm{Ap}(S, n_i)$ and $n - n_i \in \mathrm{Ap}(S, n_j)$.

- If either i or j, say i, is 0 then $n - n_j = \lceil\frac{k}{x}\rceil ha + kd$ for some $0 \le k < a$ by Theorem 3.2. Then

$$\begin{aligned}
\mathrm{t}(n) \le \max \mathsf{L}(n) &= \max \mathsf{L}\big((\lceil\frac{k}{x}\rceil h + h)a + (k+j)d\big) \\
&= \lceil\frac{k}{x}\rceil h + h + \lfloor\frac{k+j}{a}\rfloor d - \lceil\frac{(k+j) \mod a}{x}\rceil(h-1) \\
&\le \lceil\frac{k}{x}\rceil h + h + \lfloor\frac{k+j}{a}\rfloor d \\
&\le \lceil\frac{a-1}{x}\rceil h + h + \lfloor\frac{k+j}{a}\rfloor d \\
&\le \lceil\frac{a-1}{x}\rceil h + h + d = t.
\end{aligned}$$

- Let neither i nor j is 0. Assume that $\mathrm{t}(S) > t$. Then either z or z' is greater than t. This implies that either $n - n_i$ or

$n - n_j$ has order greater than or equal to t. Let for example $\max \mathsf{L}(n - n_i) \geq t$. Since $n - n_i - n_j \notin S$ we have

$$\lceil \frac{(k-i-j) \mod a}{x} \rceil h > q - 2h + \lfloor \frac{k-i-j}{a} \rfloor d$$

and therefore

$$\begin{aligned}
\max \mathsf{L}(n - n_i) &= \max \mathsf{L}((q-h)a + (k-i)d) \\
&= \max \mathsf{L}((q - h + \lfloor \frac{k-i}{a} \rfloor d)a + ((k-i) \mod a)d) \\
&= q - h + \lfloor \frac{k-i}{a} \rfloor d - \lceil \frac{(k-i) \mod a}{x} \rceil (h-1) \\
&\leq q - h + d \\
&< \lceil \frac{(k-i-j) \mod a}{x} \rceil h + h - \lfloor \frac{k-i-j}{a} \rfloor d + d \\
&< \lceil \frac{a-1}{x} \rceil h + h + d = t,
\end{aligned}$$

a contradiction.

Therefore $\mathrm{t}(S) = t$.

We know that $\omega(S) \leq \mathrm{t}(S)$. We complete the proof by showing that $\omega(n, n_1) \geq t$, where $n = ta$. We saw that $n_1 \leq_S n$; again we consider the decomposition

$$n = \underbrace{a + \ldots + a}_{t-\text{times}}.$$

Then $n - n_1 \in \mathrm{Ap}(S, n_0)$; so $n_1 \not\leq_S n - n_0$, $\omega(n, n_1) \geq t$, and therefore $\omega(S, n) \geq t$. $\square$

Remark 3.11. In [[12], Theorem 4.6] it has been shown that in a large class of Krull monoids, the tame degree coincides with the τ invariant, whereas in numerical monoids generated by generalized arithmetic sequences it coincides with the ω invariant.

Remark 3.12. Let $S = \langle a, a+d, \ldots, a+xd \rangle$. In [4] the catenary degree and the tame degree of S are found as

$$\mathrm{c}(S) = \lceil \frac{a}{x} \rceil + d, \quad \mathrm{t}(S) = \begin{cases} \lceil \frac{a}{x} \rceil + d & \text{if } a \mod x = 1, \\ \lceil \frac{a}{x} \rceil + d + 1 & \text{otherwise.} \end{cases}$$

Since

$$\lceil \frac{a-1}{x} \rceil + 1 = \begin{cases} \lceil \frac{a}{x} \rceil & \text{if } a \mod x = 1, \\ \lceil \frac{a}{x} \rceil + 1 & \text{otherwise,} \end{cases}$$

our results for $h = 1$ agree with these results.

Remark 3.13. Let $S = \langle a, ha+d, \ldots, ha+xd\rangle$ where a, d, h, and x are positive integers, $\gcd(a,d) = 1$, and $1 \le x \le a-1$. Then

$$\mathrm{t}(S) - \mathrm{c}(S) = \big(\lceil\frac{a-1}{x}\rceil + 1 - \lceil\frac{a}{x}\rceil\big)h.$$

If $(a \mod x) \neq 1$ then $\lceil\frac{a-1}{x}\rceil + 1 - \lceil\frac{a}{x}\rceil = 1$ and therefore, even if the number of minimal generators is fixed, $\mathrm{t}(S) - \mathrm{c}(S)$ can be arbitrary large as h tends to infinity. The fact the difference of the tame degree and the catenary degree is growing holds for Krull monoids with cyclic class group or whose class group is an elementary 2-group, and every class contains a prime divisor (in both cases the difference grows with the Davenport constant). These follow from Proposition 6.5.2(2), Theorem 6.5.3(1), and Theorem 6.4.2 in [10].

Remark 3.14. Let $S = \langle a, ha+d, \ldots, ha+xd\rangle$ where a, d, h, and x are positive integers, $\gcd(a,d) = 1$, and $1 \le x \le a-1$. It is known that [[14], Theorem 2.8]

$$g(S) = \lceil\frac{a-1}{x}\rceil ha + ad - a - d.$$

Let

$$B = \frac{g(S) + n_p}{n_1} + 1 = (\lceil\frac{a-1}{x}\rceil + 1)h + \frac{x-1}{a}d.$$

Then

$$B - \mathrm{t}(S) = \frac{x-1}{a}d$$

and

$$B - \mathrm{c}(S) = (B - \mathrm{t}(S)) + (\mathrm{t}(S) - \mathrm{c}(S)) = \big(\lceil\frac{a-1}{x}\rceil + 1 - \lceil\frac{a}{x}\rceil\big)h + \frac{x-1}{a}d.$$

Firstly let $(a \mod x) = 1$. Then $\mathrm{c}(S) = \mathrm{t}(S)$ while $B - \mathrm{c}(S) = (x-1)d/a$ can be arbitrary large as we let d goes to infinity.

Secondly let $(a \mod x) \neq 1$ and choose d so that $(x-1)d < a$. Then $\lfloor B\rfloor = \mathrm{t}(S)$ while $B - \mathrm{c}(S) = h + (x-1)d/a$ can be arbitrary large as we let h goes to infinity.

ACKNOWLEDGMENT

We thank A. Geroldinger, P. A. García-Sánchez and S. T. Chapman for their help in developing this paper.

REFERENCES

[1] J. Amos, S. T. Chapman, N. Hine and J. Paixao, Sets of lengths do not characterize numerical monoids, Integers 7(2007), #A50

[2] D. F. Anderson, S. T. Chapman, N. Kaplan, D. Torkornoo, An Algorithm to compute ω-primality in a numerical monoid, preprint.

[3] C. Bowles, S. T. Chapman, N. Kaplan, D. Reiser, On Delta sets of numerical monoids, J. Algebra Appl., 5 (2006) 1–24.

[4] S. T. Chapman,P. A. García-Sánchez, D. Llena, The catenary and tame degree of numerical semigroups, Forum Math. 21(2009), 117–129.

[5] S. T. Chapman, P. A. García-Sánchez, D. Llena, V. Ponomarenko, J. C. Rosales, The catenary and tame degree in finitely generated commutative cancellaive monoids, Manuscripta Math.,120 (2006) 253–264.

[6] S. T. Chapman, P. A. García-Sánchez, D. Llena, J. C. Rosales, Presentations of finitely generated cancellative commutative monoids and nonnegative solutions of systems of linear equations, Discrete Appl. Math., 154 (14) (2006) 1947-1959.

[7] S. T. Chapman, M. T. Holden, T. A. Moore, Full elasticity in atomic monoids and integral domains, Rocky Mountain J. Math., 36 (2006) 1437–1455.

[8] M. Delgado, P. A. García-Sánchez, J. Morais,"Numericalsgps": a `GAP` [18] package on numerical semigroups. (`http://www.gap-system.org/Packages/numericalsgps.html`).

[9] A. Geroldinger, D. J. Grynkiewicz, W. A. Schmid, The catenary degree of Krull monoids I, To appear.

[10] A. Geroldinger, F. Halter-Koch, Non-unique facorization: Algebraic, Combinatorial and Analytic Theory, vol. 278, Chapman & Hall/CRC, 2006.

[11] A. Geroldinger, F. Halter-Koch, Non-unique facorization : a survey, Multiplicative Ideal Theory in Commutative Algebra, Springer, 2006, 207-226.

[12] A. Geroldinger, W. Hassler, Local tameness of v-notherian monoids, J. Pure Appl. Algebra 212 (2008) 1509-1524.

[13] G. L. Matthews, On numerical semigroups generated by generalized arithmetic sequences, Communications in Algebra 32(9) (2004), 3459-3469.

[14] M. Omidali, F. Rahmati, On the type and minimal presentation of certain numerical semigroups, Communications in Algebra 37(4) (2009), 1275-1283 .

[15] J.C. Rosales, P.A. García-Sánchez, Numerical Semigroups, Developments in Mathematics, Vol. 20, Springer, 2010.

[16] Ö. J. Rödseth, On a linear diophantine problem of Frobenius II, J. Reine Angew. Math. 307/308 (1979) 431-440.

[17] E. S. Selmer, On the linear diophantine problem of Frobenius, J. Reine Angew. Math. 293/294 (1977) 1-17.

[18] The `GAP` Group, Gap – Groups, Algorithms, and Programming, Version 4.4, 2004. (`http://www.gap-system.org`).

DEPARTMENT OF MATHEMATICS, BU-ALI SINA UNIVERSITY, HAMEDAN, IRAN
E-mail address: `mehdioa@gmail.com`